

HIV Transmission in Male Serodiscordant Couples in Australia, Thailand and Brazil

Andrew E Grulich,¹ Benjamin R Bavinton,¹ Fengyi Jin,¹ Garrett Prestage,^{1,2} Iryna Zablotska,¹ Beatriz Grinsztejn,³ Nittaya Phanuphak,⁴ Richard Moore,⁵ and Kersten K Koelsch,¹ on behalf of the Opposites Attract Study Group

¹ Kirby Institute, University of New South Wales, Sydney, Australia. ² Australian Research Centre in Sex Health and Society, La Trobe University, Melbourne, Australia.

³ Instituto de Pesquisa Clínica Evandro Chagas, Fundação Oswaldo Cruz, Rio de Janeiro, Brazil. ⁴ Thai Red Cross AIDS Research Centre, Bangkok, Thailand. ⁵ Northside Clinic, Melbourne, Australia.

Background

Numerous prospective studies have demonstrated that HIV transmission is greatly reduced in heterosexual HIV serodiscordant couples when the HIV-positive partner is receiving combination anti-retroviral therapy (cART) and has undetectable viral load. Comparable data in homosexual male serodiscordant couples are extremely limited.

We report a pre-specified interim analysis of the relationship between undetectable viral load and HIV transmission in the ongoing *Opposites Attract* observational cohort study of homosexual male serodiscordant couples from three cities in Australia (Sydney, Melbourne, and Brisbane), in Bangkok, Thailand, and in Rio de Janeiro, Brazil.

Results

By December 2014, 234 homosexual male serodiscordant couples were enrolled: 135 from Australia, 52 from Bangkok and 47 from Rio de Janeiro. Median age was 36.1 years for HIV-negative partners and 35.9 years for HIV-positive partners. Over one-third of couples (38.7%) had been having sex with each other for less than 12 months, while 32.9% and 28.4% had been having sex with each other for 1-5 years and for over 5 years, respectively.

There were a total of 150.0 CYFU in 152 couples with at least one follow-up visit of whom 65 (42.8%) were in a non-monogamous relationship. At baseline, 84.2% of HIV-positive partners were on cART and in total 82.9% had undetectable viral load. STI prevalence was 11.2% in HIV-positive partners and 6.6% in HIV-negative partners.

There were 90.8 CYFU in periods where CLAI was reported with a total of 5,905 acts of CLAI in 88 couples. There were no linked HIV transmissions. The upper limit of the 95% confidence interval of the transmission rate was 4.06 per 100 CYFU for periods in which CLAI was reported, and 6.46 per 100 CYFU for periods in which receptive CLAI was reported.

Conclusions

There were no linked HIV transmissions in 150 CYFU in these homosexual male serodiscordant couples, despite close to six thousand acts of CLAI. The upper confidence limit of the transmission rate during follow-up in periods during which CLAI was occurring was 4.06 per 100 CYFU. These data add to emerging evidence that the rate of HIV transmission in homosexual male serodiscordant couples is very low when the HIV-positive partner is on cART and has undetectable viral load. Further follow-up of a larger sample size is required to accurately delineate any residual risk, and the *Opposites Attract Study* continues to recruit and follow-up homosexual male serodiscordant couples.

Method

Homosexual male serodiscordant couples reporting regular anal intercourse with each other were recruited through clinical sites. Detailed information on sexual risk behaviours was collected at each visit from the HIV-negative partner. HIV-negative partners were tested at baseline and follow-up for HIV antibodies and sexually transmitted infections (STIs), and HIV-positive partners were tested for HIV viral load and STIs. HIV incidence rates were calculated per couple-year of follow-up (CYFU) using person-year methods, and stratified by whether different forms of condomless anal intercourse (CLAI) were reported. Undetectable viral load was defined as less than 200 copies per millilitre. One-sided confidence intervals were calculated using the exact Poisson method. Linked HIV transmission in couples was defined by phylogenetic analysis.

Table 1: HIV incidence by category of condomless anal intercourse reported

Type of intercourse without a condom reported by HIV-negative partner	Linked trans-missions (n)	Couple-years of follow up (CYFU)	No. of CLAI ^a acts	Incidence rate per 100 CYFU (95% CI)
Overall	0	149.96	5,905	0 (0-2.46)
Any CLAI	0	90.83	5,905	0 (0-4.06)
Insertive CLAI	0	77.87	3,569	0 (0-4.74)
Receptive CLAI	0	57.08	2,337	0 (0-6.46)
Any CLAI when VL ^b <200 copies	0	88.59	5,656	0 (0-4.16)
Any CLAI when VL >200 copies	0	2.00	237	0 (0-184.31)

^a CLAI refers to condomless anal intercourse. ^b VL refers to viral load

Acknowledgements

The Kirby Institute receives funding from the Australian Government Department of Health and Ageing. The Opposites Attract Study receives funding from the Australian National Health and Medical Research Council (NHMRC) and the Foundation for AIDS Research (amfAR). No pharmaceutical grants were received in the development of this study. The Opposites Attract Study Group also includes: David Baker, Mark Bloch, David A Cooper, Nicholas Doong, Sean Emery, Christopher Fairley, Jennifer Hoy, Anthony Kelleher, Anna McNulty, Kate Merlin, David Orth, Catherine Pell, Norm Roth, Darren Russell, BK Tee, David Templeton, Kathy Triffitt, David Wilson. Thanks to the participants in the study, clinical staff in the participating clinics, laboratory staff at participating laboratories, the funders, community partner organisations, Lara Cassar (research assistant), and Angie Pinto (phylogenetic analysis).

Presented at the Conference on Retroviruses and Opportunistic Infections (CROI 2015), Washington State Convention Center, Seattle, Washington. 23 to 26 February 2015. Poster Number: 1019LB