









Voluntary Medical Male Circumcision as a Platform for Non-Communicable Disease Case Finding in Namibia

by: Nikki Soboil¹, Shawn Rooinasie¹, Catharine Laube¹, Abubakari Mwinyi¹, and Amir Shaker² affiliation: ¹Jhpiego; ²Ministry of Health and Social Services, Erongo Directorate, Republic of Namibia

Background

- The burden of non-communicable diseases (NCDs), including hypertension (HTN), is growing in sub-Saharan Africa, particularly in urban areas, with evidence of considerable underdiagnosis.1
- A systematic review of HTN in sub-Saharan Africa found that less than 40% of people with HTN had been previously diagnosed.²
- HTN is more prevalent in African males and increases with age.²
- Males, who seek health care less than females, are particularly likely to suffer from undiagnosed HTN and other NCDs.
- In Namibia, it is estimated that 38% of the urban population are living with HTN.³
- Voluntary medical male circumcision (VMMC) is one of few preventive health services catering specifically to males.
- Optimal VMMC service delivery infrastructures provide ideal health screening opportunities, including NCD screening.
- The Namibian Ministry of Health and Social Services and Jhpiego launched high-volume, nurse-led VMMC services at Swakopmund State Hospital in Erongo Region, Namibia, in May 2016.

Namibja

wakopmund

Over 90% of the clients served through July 31, 2016, were aged 20 years or older, in contrast to VMMC clients across eastern and southern Africa, the majority of whom have been aged between 10 and 19 years.4

HTN Screening

- All clients received full physical screening prior to VMMC services, including preoperative screening for HTN, diabetes, bleeding disorders, and TB.
- Blood pressure (BP) was measured using the appropriate cuff size and taken at the level of the
- Clients presenting with BP at or above 140/90 mm Hg were remeasured after progressively widening intervals of rest (1, 15, and 30 minutes), as needed.
- If BP remained elevated after four measurements, clients were referred to the Outpatient Department for review and management, and surgery was deferred.

- Jhpiego abstracted data from client records for males registered for VMMC services between May 13 and July 31, 2016, including preoperative physical screening data, to characterize the proportion of clients with BP at or above 140/90 mmHg.
- A random sample of 28 suspected hypertensive clients were contacted post hoc to determine whether they had been previously diagnosed.

Classification of HTN

Table 1. HTN classification according to elevated systolic or diastolic BP (mmHg)

Classification	Systolic BP (mmHg)	Diastolic BP (mmHg)
Normal to high normal (pre) HTN	120–139	80–89
Mild HTN—stage 1	140–159	90–99
Moderate to severe HTN—stage 2	> 160	> 100

Source: Ministry of Health and Social Services [Namibia]. 2011. Namibia Standard Treatment Guidelines. http://apps.who.int/medicinedocs/documents/s19260en/s19260en.pdf.

Results

Figure 1. Screened Clients

Of the 1,266 males screened for VMMC between May 13 and July 31, 2016, 367 (29%) were suspected to suffer from HTN due to a BP reading at or above 140/90 mmHg.




Figure 2. Hypertensive

- Classification of clients suspected to have HTN according to average BP measurements (n = 367)
- **269 (73%) stage 1**





Figure 3. Isolated Systolic Hypertension

- Classification of clients suspected to have ISH (> 140/< 90 mmHg) according to average BP measurements (n = 142)
- **135 (95%) stage 1**
- 6 (4%) stage 2
- 1 (1%) stage 3



Classification of Isolated Systolic Hypertension (ISH)

Table 2. ISH classification according to elevated systolic

Classification	Systolic BP (mmHg)	Diastolic BP (mmHg)
Stage 1	140–159	< 90
Stage 2	160–179	< 90
Stage 3	> 180	< 90

Source: Whitworth JA; World Health Organization, International Society of Hypertension Writing Group. 2003. 2003 World Health Organization (WHO)/International Society of Hypertension (ISH) statement on management of hypertension. *J Hypertens*. 21(11):1983–1992. doi: 10.1097/01.hjh.0000084751.37215.d2.

Table 3. Age and HTN screening outcome of 1,266 VMMC clients screened May 13-July 31, 2016

	Age in Years							Total	
	15–19	20–24	25–29	30–34	35–39	40–44	45–49	50+	IOlai
Number of clients with normal BP	54	243	252	162	90	63	18	17	899
Number of HTN suspects	15	73	93	71	41	40	14	20	367
Total number screened	69	316	345	233	131	103	32	37	1,266
Percentage of clients that were HTN suspects	22	23	27	30	31	39	44	54	29

95% of clients suspected to suffer from HTN (n = 367) were aged 15–49 years (young

	Age in Years							Total	
	15–19	20–24	25–29	30–34	35–39	40–44	45–49	50+	Total
Stage 1	11	62	75	49	30	26	7	9	269
Stage 2	4	11	18	22	11	14	7	11	98
Total	15	73	93	71	41	40	14	20	367

Table 4. Age and classification of VMMC clients screened and suspected to suffer	
from HTN (n = 367)	

adults < 50 years of age)

Table 5. Age breakdown of random sample (n = 28) contacted to determine HTN

Of the random sample of 28 suspected hypertensive clients contacted post hoc:

- 13 (46%) had been diagnosed with HTN prior to VMMC
- 15 (54%) were newly diagnosed with HTN as result of VMMC screening
- 14 (93%) of those newly diagnosed were aged < 50 years

status prior to VMMC screening

	Age in Years						Takal		
	15–19	20–24	25–29	30–34	35–39	40–44	45–49	50+	Total
Number (%) known	0	0	2 (7)	0	2 (7)	1 (4)	4 (14)	4 (14)	13 (46)
Number (%) unknown	1 (4)	0	4 (14)	7 (25)	1 (4)	1 (4)	0	1 (4)	15 (54)
Total	1	0	6	7	3	2	4	5	28

Conclusions

- VMMC screening represents an important opportunity for HTN and other non-communicable disease (NCD) case finding, particularly in programs serving male clients aged 15-49 in geographic areas where considerable underdiagnosis is evident.
- Individuals suffering from HTN at an early age are at risk for developing various health problems, including hardening of the arteries, stroke, brain hemorrhage, kidney malfunction, and blindness. In addition to being at increased risk, they also experience higher rates of death from stroke and kidney disease than the general population. Increased early HTN diagnosis is shown to improve males' management of their cardiovascular health, thus case finding in VMMC services has the potential to contribute to reduced morbidity and reduced burden to the health care system from advanced cardiovascular disease and treatment.
- VMMC programs seeking to attract a greater proportion of males aged 15–29 should prioritize careful preoperative physical screening, as well as a systematic approach to deferrals and active referrals for clients diagnosed with HTN.
- Service delivery models integrated/co-located with primary care may help reduce loss to follow-up for males newly diagnosed with HTN. Research is needed to better understand the full NCD disease burden in VMMC clients within and outside of Namibia.

References

1. Guariguata L, de Beer I, Hough R, Mulongeni P, Feeley FG, Rinke de Wit TF. 2015. Prevalence and knowledge assessment of HIV and non-communicable disease risk factors among formal sector employees in Namibia. *PLoS ONE*. 10(7):e0131737. doi:10.1371/ journal.pone.0131737.

2. Addo J, Smeeth L, Leon D. 2007. Hypertension in sub-Saharan Africa. *Hypertension*. 50:1012–1018.

3. Hendriks ME, Wit FWNM, Roos MTL, et al. 2012. Hypertension in sub-Saharan Africa: cross-sectional surveys in four rural and urban communities. PLoS ONE. 7(3):e32638. doi:10.1371/journal.pone.0032638.

4. Kripke K, Njeuhmeli E, Samuelson J, et al. 2016. Assessing progress, impact, and next steps in rolling out voluntary medical male circumcision for HIV prevention in 14 priority countries in Eastern and Southern Africa through 2014. *PLoS ONE*. 11(7):e0158767. doi: 10.1371/journal.pone.0158767.

