



The Impact of Weight Gain and Sex on Immune Activation Following Initiation of ART

Sara H. Bares¹, Laura M. Smeaton², Vincent Vu², Sarah E. Scott³, Beth A. Smith³, Catherine Godfrey⁴, Grace A. McComsey³

¹University of Nebraska Medical Center, Omaha, NE, ²Harvard T.H. Chan School of Public Health, Boston, MA, ⁴Division of AIDS, NIAID, Bethesda, MD, ³University Hospitals Cleveland Medical Center and Case Western Reserve University School of Medicine, Cleveland, OH

Background

- Immune activation persists despite suppressive antiretroviral therapy (ART) and may also differ by sex or body composition.
- We hypothesized that weight gain, sex and BMI would be associated with changes in immune activation following ART initiation and explored these relationships in a selected subset of women and men who had initiated ART in two large randomized ACTG trials (A5202 and A5257).

Objectives

- To test if baseline biomarker levels differ by sex, baseline BMI category or whether person later gained weight or had stable weight over time, if these differences vary by levels of the other 2 factors, and whether these differences persist after adjustment by age, or pre-ART CD4 and viral load.
- To test if changes from baseline levels in selected inflammatory markers differ by whether person later gained weight or had stable weight over time, and explore if these differences persist after adjustment for sex, age, pre-ART BMI category, CD4 and viral load.

Methods




- A purposeful sampling design selected participants who achieved virologic suppression on ART and either maintained weight within +/- 0.5 kg/m2 ("maintainers") or gained 2.6-6.4 kg/m2 ("gainers") from baseline to 96 weeks.
- From stored plasma samples, we measured IL-6, sTNF-RI and II, IP-10, hsCRP, sCD14 and sCD163 at weeks 0 and 96.
- Multivariable linear regression explored associations of weight gain (gainers vs maintainers), sex, and pre-ART BMI with outcomes of pre-ART biomarker concentrations and changes from baseline to week 96, adjusting for baseline age, race/ethnicity, ART regimen, CD4 count and HIV-1 RNA.

Results

	Women and men who gained weight (2.6-6.4 kg/m ²) (n=136)	Women and men who maintained weight (+/- 0.5 kg/m ²) (n=144)	P-Value (weight gain)	Women (n=166)			Men (n=174)		
				Gained weight (n=96)	Maintained weight (n=70)	P-Value (weight gain within women)	Gained weight (n=100)	Maintained weight (n=74)	P-value (weight gain within men)
Clinical Trial, n (%)									
A5202	82 (42%)	55 (38%)	NA*	42 (44%)	25 (36%)	NA*	40 (40%)	30 (41%)	NA*
A5257	114 (58%)	89 (62%)		54 (56%)	45 (64%)		60 (60%)	44 (59%)	
ART Regimen, n (%)									
ATV/r + NUCs*	74 (38%)	58 (40%)	NA*	34 (35%)	23 (33%)	NA*	40 (40%)	35 (47%)	NA*
DRV/r + NUCs*	37 (19%)	29 (20%)		20 (21%)	17 (24%)		17 (17%)	12 (16%)	
EFV + NUCs*	43 (22%)	26 (18%)		26 (27%)	13 (19%)		17 (17%)	13 (18%)	
RAL + NUCs*	42 (21%)	31 (22%)		16 (17%)	17 (24%)		26 (26%)	14 (19%)	
Age (years), median (Q1, Q3)	42 (34, 49)	42 (33, 49)	NA*	43 (35, 49)	42 (34, 49)	NA*	42 (33, 49)	42 (31, 49)	NA*
Race/Ethnicity, n (%)									
White Non-Hispanic	56 (29%)	55 (38%)	0.26	13 (14%)	16 (23%)	0.46	43 (43%)	39 (53%)	0.55
Black Non-Hispanic	89 (45%)	53 (37%)		56 (58%)	35 (50%)		33 (33%)	18 (24%)	
Hispanic	48 (24%)	33 (23%)		26 (27%)	18 (26%)		22 (22%)	15 (20%)	
Other	3 (2%)	3 (2%)		1 (1%)	1 (1%)		2 (2%)	2 (3%)	
Pre-ART BMI (kg/m ²), median (Q1, Q3)	25.4 (22.4, 29.5)	25.9 (22.4, 29.0)	0.70	27.1 (23.3, 32.8)	27.7 (23.6, 32.3)	0.69	24.1 (21.7, 27.6)	24.2 (22.4, 27.1)	0.74
Pre-ART CD4 (cells/mm ³), median (Q1, Q3)	195 (47, 337)	344 (232, 460)	<.01	219 (55, 363)	328 (231, 441)	<.01	174 (45, 299)	361 (242, 465)	<.01

* Clinical trial, ART regimen, and Age used in sampling design to enforce balance by these factors; therefore, no hypothesis testing performed.
*NUCS included TDF/FTC, or ABC/3TC (only with EFV or ATV/r)

Results



Results

- While pre-ART BMI was similar between gainers and maintainers (overall and within sex), gainers had significantly lower pre-ART CD4 vs maintainers.
- In adjusted models among those with normal pre-ART BMI, pre-ART IL-6, sTNF-RII, IP-10, and sCD163 were higher for gainers versus maintainers.
- Association of weight gain on week 96 changes of these 4 biomarkers differed by sex; women who gained weight had smaller declines in biomarkers compared to men who gained.
- For sTNF-RII and IL-6, the association between weight gain and changes in these biomarkers also varied by pre-ART BMI.

- NOTE: Similar changes were noted for the other biomarkers

	Pre-ART concentrations Adjusted differences (95% CI)	Women (n=166)		Men (n=174)		Sex by weight gain interaction p-values
		Gained weight (n=96)	Maintained weight (n=70)	Gained weight (n=100)	Maintained weight (n=74)	
		Changes from baseline by sex [model based means]				
IL-6 (log ₁₀ pg/mL)	0.26 (0.17, 0.35)	-0.11 normal 0.02 overweight 0.08 obese	-0.16 -0.03 0.03	-0.25 -0.13 -0.07	-0.12 0.01 0.07	.004 ^B
sCD163 (log ₁₀ ng/mL)	0.08 (0.02, 0.14)	-0.17	-0.25	-0.23	-0.21	<.001
sTNF-RII (log ₁₀ pg/mL)	0.12 (0.07, 0.16)	-0.26 normal -0.20 overweight -0.18 obese	-0.24 -0.25 -0.25	-0.30 -0.24 -0.22	-0.21 -0.22 -0.22	.021
IP-10 (log ₁₀ pg/mL)	0.19 (0.11, 0.27)	-0.47	-0.50	-0.53	-0.41	.004

*Pre-ART concentrations for subgroup of participants with normal pre-ART BMI

Conclusions

- Among non-obese persons, higher pre-treatment immune activation markers are significantly associated with weight gain following ART initiation, even after controlling for pre-ART CD4 cell count.
- Weight gain attenuates the decline in several immune activation markers following ART initiation among women; thus, women may be at increased risk for complications of weight gain.
- Identifying individuals at risk of weight gain may allow for targeted investigation of preventive interventions.

Acknowledgements

We would like to thank the study coordinators and participants at each of the study sites for their contributions to this work. This work was funded by National Institutes of Health grants UM1 AI068636 (ACTG) and UM1 AI06834 (SDAC) as well as an Office for Research on Women's Health supplement to 1U01 AI069501.