Sexual risk and HIV testing disconnect in MSM recruited to an HIV self-testing trial

Alison Rodger¹, David Dunn¹, Andrew N. Phillips¹, Leanne McCabe¹, Peter Weatherburn², Fiona Lampe¹, T Charles Witzel², Fiona Burns¹, Denise Ward¹, Roger Pebody³, Roy Trevelion⁴, Peter Kirwan⁵, Yolanda Collaco-Moraes¹, Sheena McCormack¹

Email: alison.rodger@ucl.ac.uk http://www.selphi.org

¹University College London, London, UK. ²London School of Hygiene & Tropical Medicine, London, UK. ⁴HIV iBase, London, UK. ⁵Public Health England, UK.

Background

Increasing levels of HIV testing in men who have sex with men (MSM) remain key to reducing incidence. However, levels of ever and repeat HIV testing in UK MSM do not meet current testing recommendations with 28% never testing and only 55% testing annually (1, 2). Particularly poorly adhered to is the recommendation to test 3 monthly for men at higher risk of HIV through recent condomless anal intercourse (CAI) with partners of unknown or serodifferent HIV status, and/or drug use during sex (chemsex) with only 27% of those at 'higher risk' of HIV infection testing even 6 monthly (1).

HIV self-testing (HIVST) may increase testing rates by removing structural and social barriers to testing (3). We report on the frequency of previous HIV testing and associated factors at baseline in MSM who opted to enroll in a large HIV self-testing RCT (SELPHI).

Methods

SELPHI is an internet based, open-label, randomised controlled trial, which aims to assess effectiveness of free HIVST kits on HIV diagnosis rates.

Criteria for enrolment were aged ≥16 years old, male (including trans man), ever had anal intercourse (AI) with a man, living in England or Wales, not known to be HIV positive and provided consent to link to national HIV surveillance databases (4). Participants were randomly allocated 3:2 at enrolment to a free HIVST versus no free HIVST (Randomisation A).

Data collected via an online survey included sociodemographics (gender, sexual identity, education, age, ethnicity, UK birth), sexual behaviour, HIV/STI testing history, pre-exposure prophylaxis (PrEP) use.

Figure 1: Trial Scheme for Randomisation A



Sociodemographic characteristics were similar in men at higher risk of HIV (defined as 2+CAI partners in the previous 3) months) to the overall sample. However PrEP use varied with 3.8% currently using PrEP in all participants compared to 7.8% men at higher risk of HIV infection were currently using PrEP.

This study is funded by the National Institute for Health Research (NIHR) under its Programme Grants for Applied Research Programme (Grant Reference Number RP-PG-1212-20006). The views expressed are those of the authors and not necessarily those of the NIHR or the Department of Health and Social Care.

Table 1. Baseline characteristics at baseline of all recruited and of men at high risk of HIV infection

aracteristic	All participants (n=10,112)	Participants at current high risk			
gion					
ondon	2510 (24.8)	938 (23.7)			
E England	1430 (14.2)	562 (14.2)			
ther	6166 (61.0)	2449 (62.0)			
stance to nearest GUM clinic		, , ,			
n)					
2	2795 (29.5)	1104 (29.6)			
-5	3737 (39.4)	1419 (38.1)			
-10	1616 (17.0)	659 (17.7)			
10	1328 (14.0)	546 (14.6)			
e (years)					
6-19	534 (5.3)	205 (5.2)			
0-29	3301 (32.6)	1299 (32.9)			
0-39	2884 (28.5)	1144 (28.9)			
0-49	1919 (19.0)	770 (19.5)			
50	1474 (14.6)	534 (13.5)			
ins gender	81 (0.8)	32 (0.8)			
nnicity					
/hite British	7595 (75.5)	3011 (76.6)			
/hite Other	1406 (14.0)	501 (21.7)			
lack	161 (1.6)	79 (2.0)			
sian	437 (4.3)	155 (3.9)			
lixed	313 (3.1)	119 (3.0)			
ther	150 (1.5)	66 (1.7)			
shest level education					
chool	4169 (41.7)	1776 (45.6)			
ollege	1119 (11.2)	465 (11.9)			
niversity	4706 (47.1)	1655 (42.4)			
. male anal sex partners in					
st 3 months					
	1064 (10.5)	-			
	2140 (21.2)	-			
-4	4082 (40.4)	1821 (46.1)			
.9	1894 (18.7)	1336 (33.8)			
10	932 (9.2)	795 (20.1)			
. male condom-less anal sex					
rthers in past 3 months					
	2828 (30.0)	-			
	3331 (32.9)				
-4	2943 (29.1)	1821 (46.1)			
9	700 (6.9)	1336 (33.8)			
	309 (3.1)	795 (20.1)			
		F40 (12 C)			
/er	795 (7.9)	540(13.6)			
	389 (3.8)	3UI (7.6)			

*2+CAI partners in previous 3 months

Results

10,112 men were recruited to the trial including 81 (0.8%) trans men. Median age was 33 years (IQR 26-44). Most were of white ethnicity (89.5%) with only 1.6% of black ethnicity and 4.3% of Asian ethnicity. The men were highly educated with 47.1% having a university education. In the previous 3 months, 89% reported AI and 72% reported CAI with ≥1 male partner (Table1).

Characteristic	All participants (n=10,112)	Participants currently at high risk of HIV infection (n=3,952)
Never tested	1537 (15.4)	449 (11.5)
Time since last HIV test (months)		
<3	1695 (16.9)	884 (22.6)
3-6	1628 (16.3)	729 (18.6)
6-12	2093 (20.9)	822 (21.0)
12-24	1765 (17.6)	648 (16.6)
>24	1297 (12.9)	382 (9.8)
Number of HIV tests in previous 12		
months (if at least one)		
1	2806 (52.3)	1116 (46.9)
2	1419 (26.8)	645 (27.1)
3	657 (12.7)	365 (15.30
≥4	402 (7.6)	253 (10.6)
Venue of last HIV test		
Sexual health clinic	5090 (61.2)	2235 (65.6)
Hospital	415 (5.0)	136 (4.0)
Community service	347 (4.2)	127 (3.7)
Self-sample	1380 (16.6)	504 (14.8)
Self-test	556 (6.7)	212 (6.2)
Bar/pub/club/sauna	114 (1.4)	58 (1.7)
GP	294 (3.5)	102 (3.0)
Other	121 (1.4)	32 (0.9)

Overall, 17%, 33%, 54%, and 72% had tested for HIV in the last 3 months, 6 months, 12 months, and 2 years respectively. 13% had tested >2 years ago and 15% had never tested. In men had higher risk of HIV: 12% had never tested and almost half (47%) had tested > 6 months previously (Table 2, Figure 2).



Considering a multivariable logistic analysis of factors associated with having never tested for HIV, living outside of major urban regions such as London and the southeast of England; being aged 16 to 20 years and having a lower level of education were all associated with never having been tested for HIV (data not shown).

- Care. 2014;26(3):297-303.
- 2015;19(11):1949-65

Table 2. HIV testing characteristics of all participants and those at high risk of HIV infection

Figure 2: Time since previous HIV test in all men and men at high risk of HIV infection

1. McDaid LM, et al. Frequency of HIV testing among gay and bisexual men in the UK: implications for HIV prevention. HIV Med.17(9):683-93 2. Knussen C, Flowers P, McDaid LM. Factors associated with recency of HIV testing amongst men residing in Scotland who have sex with men. AIDS

3. Figueroa C, Johnson C, Verster A, Baggaley R. Attitudes and Acceptability on HIV Self-testing Among Key Populations: A Literature Review. AIDS Behav.

4. Gabriel MM, et al. Protocol, rationale and design of SELPHI: a randomised controlled trial assessing whether offering free HIV self-testing via the internet increases the rate of HIV diagnosis.BMC Infect Dis. 2018. 18(1):531

Characteristic	Tested at least once in in previous 6 months			
	No. (%)	Adjusted OR (95% CI)	P value	
Region			<0.001	
London	522 (56.2)	1.95 (1.64,2.31)		
SE England	253 (45.3)	1.55 (1.28,1.88)		
Other	837 (34.5)	1.00		
Distance to nearest GUM clinic			0.02	
(km)				
<2	537 (49.0)	1.20 (1.01,1.42)		
2-5	592 (42.2)	1.00		
5-10	224 (34.2)	0.86 (0.71,1.06)		
≥10	195 (36.1)	0.97 (0.78,1.21)		
Age (years)			0.11	
16-19	61 (29.9)	0.74 (0.53,1.04)		
20-29	547 (42.3)	1.00		
30-39	501 (44.1)	0.96 (0.81,1.13)		
40-49	314 (41.6)	0.90 (0.75,1.09)		
≥50	190 (36.0)	0.77 (9.62,0.96)		
Ethnicity			0.01	
White British	1134 (38.0)	1.00		
White Other	270 (54.7)	1.36 (1.11,1.68)		
Black	38 (48.1)	1.18 (0.74,1.88)		
Asian	71 (46.1)	0.84 (0.60,1.19)		
Mixed	48 (40.3)	0.86 (0.58,1.26)		
Other	39 (60.0)	1.57 (0.93,2.66)		
Highest level education			<0.001	
School	570 (32.4)	0.55 (0.47,0.63)		
College	185 (40.2)	0.71 (0.57,0.88)		
University	834 (50.8)	1.00		

In multivariate logistic regression of selected sociodemographic factors in men at higher risk of HIV, region of residence and education level were most strongly associated with a higher likelihood of having a recent HIV test (Table 3).

Conclusions

MSM recruited to the SELPHI HIVST RCT were not testing in line with current UK recommendations. Overall, almost half had never tested or last tested >12 months previously. In men at higher risk of recent HIV, less than half had tested in previous 6 months and a tenth had never tested. This may be due to low perceived risk of HIV infection, structural barriers (e.g. clinics difficult to access because of time constraints or capacity issues or distance) and individual psycho-social issues including perceived stigma.

Associations with lower levels of ever and repeat testing in this study suggest that targeting efforts and expanding testing opportunities in younger men and in those with lower levels of education or who live further from testing centres may improve access and uptake of HIV testing. Online promotion of free HIVST may be key to addressing many of these barriers. However we recruited low levels of BME and trans men who are key MSM groups at risk of HIV infection.

Further work is required to ensure that HIVST service provision can be adapted and targeted more effectively meet the needs of these key populations.





Table 3. Logistic regression of predictors of recent testing among men at high risk of HIV infection





Smarter studie Global impact Better health

