

Lameck Chinula¹, Evan Goldberg², Lauren Ziemba², Katie McCarthy³, Chelsea Krotje⁴, Kevin Knowles⁴, Jyoti Mathad⁵, Maria Leticia Santos Cruz⁶, Risa Hoffman⁷, Patrick Jean-Philippe⁸, Tsungai Chipato⁹, Deo Wabwire¹⁰, Judith Currier⁷, Shahin Lockman^{11,12}, Nahida Chakhtoura¹³ for the IMPAACT 2010/VESTED Study Team and Investigators

¹Division of Global Women's Health, Department of Obstetrics and Gynecology, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA; ²UNC Project Malawi, Lilongwe, Malawi; ³Center for Biostatistics in AIDS Research, Harvard TH Chan School of Public Health, Boston, MA, USA; ⁴FHI 360, Durham, NC, USA; ⁵Frontier Science Foundation, Amherst, NY, USA; ⁶Department of Medicine, Weill Cornell Medical College, New York, NY, USA; ⁷Hospital Federal dos Servidores do Estado, Rio de Janeiro, Brazil; ⁸David Geffen School of Medicine, Division of Infectious Diseases, University of California, Los Angeles, CA, USA; ⁹National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD, USA; ¹⁰College of Health Sciences, University of Zimbabwe, Harare, Zimbabwe; ¹¹Makerere University - Johns Hopkins University Research Collaboration, Kampala, Uganda; ¹²Division of Infectious Disease, Brigham and Women's Hospital, Boston, MA, USA; ¹³Department of Immunology and Infectious Diseases, Harvard TH Chan School of Public Health, Boston, MA, USA; ¹⁴Botswana Harvard AIDS Institute Partnership, Gaborone, Botswana; ¹⁵Eunice Kennedy Shriver National Institute of Child Health & Human Development, Bethesda, MD, USA.

BACKGROUND

In IMPAACT 2010, a trial comparing 3 antiretroviral treatment (ART) regimens initiated in pregnancy, found:


- Dolutegravir (DTG), particularly in combination with emtricitabine (FTC) and tenofovir alafenamide (TAF), was associated with more pregnancy weight gain, though lower than the Institute of Medicine's recommended weekly weight gain in 2nd and 3rd trimester
- DTG+FTC/TAF had better pregnancy outcomes than ART containing FTC with efavirenz (EFV) or tenofovir disoproxil fumarate (TDF)

The impact of these weight gain patterns during pregnancy on gestational diabetes is unknown. We evaluated glycated hemoglobin (HbA1c) and glucose levels and their association with maternal ART regimen in a subset of IMPAACT 2010 participants.

METHODS

- 643 pregnant women with HIV in 9 countries were randomized 1:1:1 to start open-label DTG+FTC/TAF, DTG+FTC/TDF, or EFV/FTC/TDF at 14-28 weeks' gestation
- Following protocol amendment, in a subset of participants, we sampled maternal HbA1c and random glucose at study entry, 12 weeks on study (antepartum) and delivery; and neonatal glucose within 48 hours of birth
- We assessed by-arm differences in mean levels and time-averaged area under the curve (AUC) of HbA1c and glucose in women (and glucose in infants) using student's t-tests
- We assessed by-arm differences in the proportions of women with post-baseline HbA1c $\geq 5.7\%$ (marker of pre-diabetes) or $\geq 6.5\%$ (marker of diabetes) using Wald Tests

FIGURE 1. Mean HbA1C (%) by randomized arm



We observed no significant differences in maternal HbA1C and no clinically meaningful differences in maternal or infant random glucose in women starting DTG+FTC/TAF, DTG+FTC/TDF, vs EFV/FTC/TDF in pregnancy or their newborns, despite differences in pregnancy weight gain among the regimens


RESULTS

- 348 mothers and 65 infants included in this analysis: DTG+FTC/TAF (114), DTG+FTC/TDF (116), EFV/FTC/TDF (118) arms
- Maternal enrolment medians were: age 25.9 years, gestational age 21.5 weeks, BMI 24.1 kg/cm², HIV-1 RNA 3.1 log₁₀, and CD4 466 cells/mm³
- Maternal mean HbA1c levels and mean time-averaged HbA1c AUC did not differ significantly between arms (Table 1, Figure 1)
- 7.0% of women in the DTG+FTC/TAF arm, 6.0% in the DTG+FTC/TDF arm, and 3.4% in the EFV/FTC/TDF arm had at least one post-baseline HbA1c $\geq 5.7\%$ (p ≥ 0.21 for between-arm comparisons) (Table 1). A similar pattern was observed across the randomized arms when women with HbA1c $\geq 5.7\%$ at baseline were excluded (Figure 2)
- No women had a diabetes diagnosis at entry; one woman in the DTG+FTC/TAF arm developed HbA1c $\geq 6.5\%$ after enrollment
- The DTG+FTC/TDF arm had slightly higher mean time-averaged AUC glucose (4.80 mmol/L) than the EFV/FTC/TDF arm (4.63 mmol/L, mean difference [95% CI]: 0.17 [0.00, 0.34] mmol/L) (Table 1)
- Infant glucose levels ≤ 48 hours of birth were similar by arm (Table 1)

TABLE 1. Maternal antepartum HbA1c and maternal and infant random glucose by randomized arm

	DTG+FTC/TAF	DTG+FTC/TDF	EFV+FTC/TDF	Treatment Arm Comparison	Difference in Means or Percentages (95% CI)	P-value
Mean (standard deviation) HbA1c at delivery (%)	5.14 (0.31)	5.11 (0.38)		DTG+FTC/TAF vs DTG+FTC/TDF	0.03 (-0.06, 0.12)	0.51
		5.11 (0.38)	5.10 (0.32)	DTG+FTC/TDF vs EFV/FTC/TDF	0.01 (-0.09, 0.10)	0.88
	5.14 (0.31)		5.10 (0.32)	DTG+FTC/TAF vs EFV/FTC/TDF	0.04 (-0.05, 0.12)	0.37
% (n/N) of women with HbA1c $\geq 5.7\%$ at either antepartum week 12 or delivery	7.0% (8/114)	6.0% (7/116)		DTG+FTC/TAF vs DTG+FTC/TDF	-1.0% (-7.9%, 5.8%)	0.76
		6.0% (7/116)	3.4% (4/118)	DTG+FTC/TDF vs EFV/FTC/TDF	-2.6% (-8.9%, 3.2%)	0.34
	7.0% (8/114)		3.4% (4/118)	DTG+FTC/TAF vs EFV/FTC/TDF	-3.6% (-10.2%, 2.4%)	0.21
Mean (sd) time-averaged maternal glucose AUC (mmol/L)	4.66 (0.72)	4.80 (0.66)		DTG+FTC/TAF vs DTG+FTC/TDF	-0.14 (-0.32, 0.04)	0.13
		4.80 (0.66)	4.63 (0.64)	DTG+FTC/TDF vs EFV/FTC/TDF	0.17 (0.00, 0.34)	0.045
	4.66 (0.72)		4.63 (0.64)	DTG+FTC/TAF vs EFV/FTC/TDF	0.03 (-0.15, 0.21)	0.72
Mean (sd) infant glucose within 48 hours of birth (mmol/L)	3.43 (1.09)	3.40 (1.06)		DTG+FTC/TAF vs DTG+FTC/TDF	0.03 (-0.63, 0.69)	0.93
		3.40 (1.06)	3.49 (1.03)	DTG+FTC/TDF vs EFV/FTC/TDF	-0.09 (-0.72, 0.53)	0.77
	3.43 (1.09)		3.49 (1.03)	DTG+FTC/TAF vs EFV/FTC/TDF	-0.06 (-0.74, 0.62)	0.86

FIGURE 2. HbA1c $\geq 5.7\%$ at Antepartum Week 12 or at Delivery by randomized arm



LIMITATIONS

- Modest sample size
- Glucose was not measured after fasting state

CONCLUSIONS

In this randomized trial comparing DTG+FTC/TAF, DTG+FTC/TDF, and EFV/FTC/TDF started in pregnancy, we observed:

- No significant differences in maternal antepartum/delivery HbA1C by ART regimen
- No clinically meaningful differences in maternal or infant random glucose

ACKNOWLEDGEMENTS

The IMPAACT 2010/VESTED Protocol Team gratefully acknowledges the dedication and commitment of the study participants, their communities, and CAB representatives, without whom this study would not have been possible. The authors also wish to acknowledge the IMPAACT 2010/VESTED Protocol team, NIAID, NICHD, and NIMH, and the twenty-two IMPAACT sites and staff. The study products were provided by ViiV Healthcare Ltd, Gilead Sciences, and Mylan.

REFERENCE

1. Lockman S, Brummel SS, Ziemba L, et al. Efficacy and safety of dolutegravir with emtricitabine and tenofovir alafenamide fumarate or tenofovir disoproxil fumarate, and efavirenz, emtricitabine, and tenofovir disoproxil fumarate HIV antiretroviral therapy regimens started in pregnancy (IMPAACT 2010/VESTED): a multicentre, open-label, randomised, controlled, phase 3 trial. *Lancet*. 2021 Apr 3;397(10281):1276-1292. doi: 10.1016/S0140-6736(21)00314-7. PMID: 33812487; PMCID: PMC8132194.